



Applicants: Jingrong Cao et al.
Application No.: 10/696,862

AMENDMENTS TO THE CLAIMS

1. (Currently amended) A compound of formula **I**:



or a pharmaceutically acceptable salt thereof, wherein:



R^1 is halogen, CN, NO_2 , or V_mR ;

Z^1 and Z^3 are each independently CR^Z ;

Z^2 is CR^1 ;

each occurrence of R^Z is independently halogen, CN, NO_2 , or $\text{U}_n\text{R}'$;

R^2 is $\text{U}_n\text{R}'$;

each occurrence of R^4 is independently halogen, CN, NO_2 , or V_mR ;

each occurrence of U or V is independently an optionally substituted C_{1-6} alkylidene chain, wherein up to two methylene units of the chain are optionally and independently replaced by -NR-, -S-, -O-, -CS-, -OCO-, -COCO-, -CONR-, -NRCO-, -NR CO_2 -,

Applicants: Jingrong Cao et al.
Application No.: 10/696,862

-SO₂NR-, -NRSO₂-, -CONRNR-, -NRCONR-, -OCONR-, -NRNR-, -NRSO₂NR-, -SO-, or -SO₂-;


m and n are each independently 0 or 1;

each occurrence of R is independently hydrogen or an optionally substituted C₁₋₆ aliphatic group; and each occurrence of R' is independently hydrogen or an optionally substituted C₁₋₆ aliphatic group, or a 3-8-membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur; ~~or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur~~; or R and R', two occurrences of R, or two occurrences of R', are taken together with the atom(s) to which they are bound to form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur;

Q¹ is -CO-;

R³ is Q²-Ar¹, wherein Q² is -(CHR⁶)_q-, where q is 1, 2, or 3,

or R² and Q¹-R³, taken together with the intervening nitrogen atom, form the cyclic

group: , where s is 1 or 2, each occurrence of Y is independently, as valency and stability permit, -CO-, -CS-, -SO₂-, -O-, -S-, -NR⁵-, or -C(R⁵)₂-, and R⁵ is U_nR';

Q³ is a bond or a C₁₋₆ alkylidene chain, wherein up to two methylene units of the chain are each optionally and independently replaced by -S-, -O-, -CS-, -CO₂-, -OCO-, -CO-, -COCO-, -CONR'-, -NR'CO-, -NR'CO₂-, -SO₂NR'-, -NR'SO₂-, -CONR'NR'-, -NR'CONR'-, -OCONR'-, -NR'NR'-, -NR'SO₂NR'-, -SO-, or -SO₂-; and wherein any carbon atom in the one or more methylene units is optionally substituted with one or two

Applicants: Jingrong Cao et al.
Application No.: 10/696,862

occurrences of R^6 , wherein each occurrence of R^6 is independently halogen, CN, NO_2 , or U_nR' , or two occurrences of R^6 , or R' and R^6 , taken together with the atoms to which they are bound, form an optionally substituted 3-6-membered cycloalkyl, heterocyclyl, aryl or heteroaryl ring; and

Ar^1 is a 5-8 membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from oxygen or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from oxygen or sulfur; wherein Ar^1 is optionally substituted with 0-5 independent occurrences of TR^7 ; wherein T is a bond or is a C_1 - C_6 alkylidene chain wherein up to two methylene units of T are optionally and independently replaced by -NR-, -S-, -O-, -CS-, -CO₂-, -OCO-, -CO-, -COCO-, -CONR-, -NRCO-, -NRCO₂-, -SO₂NR-, -NRSO₂-, -CONRNR-, -NRCONR-, -OCONR-, -NRNR-, -NRSO₂NR-, -SO-, or -SO₂-;

Ar^2 is a 5-8 membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur; wherein Ar^2 is optionally substituted with 0-5 independent occurrences of TR^7 ; wherein T is a bond or is a C_1 - C_6 alkylidene chain wherein up to two methylene units of T are optionally and independently replaced by -NR-, -S-, -O-, -CS-, -CO₂-, -OCO-, -CO-, -COCO-, -CONR-, -NRCO-, -NRCO₂-, -SO₂NR-, -NRSO₂-, -CONRNR-, -NRCONR-, -OCONR-, -NRNR-, -NRSO₂NR-, -SO-, or -SO₂-;

each occurrence of R^7 is independently R' , halogen, NO_2 , or CN;

each of the optional substituents of said aryl or heteroaryl ring is selected from halogen; $-R^\circ$; $-OR^\circ$; $-SR^\circ$; phenyl optionally substituted with R° ; $-O(\text{phenyl})$, optionally substituted with R° ; $-(CH_2)_{1-2}(\text{phenyl})$, optionally substituted with R° ; $-CH=CH(\text{phenyl})$,


Applicants: Jingrong Cao et al.
Application No.: 10/696,862

optionally substituted with R° ; $-\text{NO}_2$; $-\text{CN}$; $-\text{N}(R^\circ)_2$; $-\text{NR}^\circ\text{C}(\text{O})\text{R}^\circ$; $-\text{NR}^\circ\text{C}(\text{S})\text{R}^\circ$; $-\text{NR}^\circ\text{C}(\text{O})\text{N}(R^\circ)_2$; $-\text{NR}^\circ\text{C}(\text{S})\text{N}(R^\circ)_2$; $-\text{NR}^\circ\text{CO}_2\text{R}^\circ$; $-\text{NR}^\circ\text{NR}^\circ\text{C}(\text{O})\text{R}^\circ$; $-\text{NR}^\circ\text{NR}^\circ\text{C}(\text{O})\text{N}(R^\circ)_2$; $-\text{NR}^\circ\text{NR}^\circ\text{CO}_2\text{R}^\circ$; $-\text{C}(\text{O})\text{C}(\text{O})\text{R}^\circ$; $-\text{C}(\text{O})\text{CH}_2\text{C}(\text{O})\text{R}^\circ$; $-\text{CO}_2\text{R}^\circ$; $-\text{C}(\text{O})\text{R}^\circ$; $-\text{C}(\text{S})\text{R}^\circ$; $-\text{C}(\text{O})\text{N}(R^\circ)_2$; $-\text{C}(\text{S})\text{N}(R^\circ)_2$; $-\text{OC}(\text{O})\text{N}(R^\circ)_2$; $-\text{OC}(\text{O})\text{R}^\circ$; $-\text{C}(\text{O})\text{N}(\text{OR}^\circ)\text{R}^\circ$; $-\text{C}(\text{NOR}^\circ)\text{R}^\circ$; $-\text{S}(\text{O})_2\text{R}^\circ$; $-\text{S}(\text{O})_3\text{R}^\circ$; $-\text{SO}_2\text{N}(R^\circ)_2$; $-\text{S}(\text{O})\text{R}^\circ$; $-\text{NR}^\circ\text{SO}_2\text{N}(R^\circ)_2$; $-\text{NR}^\circ\text{SO}_2\text{R}^\circ$; $-\text{N}(\text{OR}^\circ)\text{R}^\circ$; $-\text{C}(=\text{NH})-\text{N}(R^\circ)_2$; $-\text{P}(\text{O})_2\text{R}^\circ$; $-\text{PO}(\text{R}^\circ)_2$; $-\text{OPO}(\text{R}^\circ)_2$; $-(\text{CH}_2)_{0-2}\text{NHC}(\text{O})\text{R}^\circ$; wherein each independent occurrence of R° is selected from hydrogen, an optionally substituted C_{1-6} aliphatic, an unsubstituted 5-6 membered heteroaryl or heterocyclic ring, phenyl, $-\text{O}(\text{phenyl})$, or $-\text{CH}_2(\text{phenyl})$, wherein optional substituents on the aliphatic group of R° are selected from NH_2 , $\text{NH}(\text{C}_{1-4}\text{aliphatic})$, $\text{N}(\text{C}_{1-4}\text{aliphatic})_2$, halogen, $\text{C}_{1-4}\text{aliphatic}$, OH , $\text{O}(\text{C}_{1-4}\text{aliphatic})$, NO_2 , CN , CO_2H , $\text{CO}_2(\text{C}_{1-4}\text{aliphatic})$, $\text{O}(\text{haloC}_{1-4}\text{aliphatic})$, or $\text{haloC}_{1-4}\text{aliphatic}$, or two independent occurrences of R° , on the same substituent or different substituents, taken together with the atom(s) to which each R° group is bound, form a 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur; and

each of the optional substituents on said alkylidene chain, aliphatic, cycloalkyl, or heterocyclyl is selected from the list of optional substituents of optional substituents for aryl and heteroaryl rings and further comprise $=\text{O}$, $=\text{S}$, $=\text{NNHR}^*$, $=\text{NN}(\text{R}^*)_2$, $=\text{NNHC}(\text{O})\text{R}^*$, $=\text{NNHCO}_2(\text{alkyl})$, $=\text{NNHSO}_2(\text{alkyl})$, or $=\text{NR}^*$, where each R^* is independently selected from hydrogen or a C_{1-6} aliphatic group;

provided that:

for compounds having the structure:



Applicants: Jingrong Cao et al.
Application No.: 10/696,862

R³ is not any one of the following groups: -CH₂(3-NHCOPh-phenyl), -CH₂-pyrrolidine, unsubstituted benzyl, -CH₂-naphthyl, -CH₂CH₂-3-(4-Cl-phenyl)-1-phenyl-1-H-pyrazol-4-yl, or -CH₂(1,3-dioxoisindole).

2-3. (Canceled)

4. (Previously presented) The compound of claim 1, wherein R² is hydrogen, or is U_nR', where n is 1, and U is a C₁₋₆ alkylidene chain wherein one or two methylene units are optionally and independently replaced by O, NR, S, or C(O).


5. (Previously presented) The compound of claim 1, wherein U is -CH₂-, -CH₂CH₂-, -CH₂CH₂CH₂-, -CH₂CH₂CH₂CH₂-, -CH₂O-, -CH₂S-, -CH₂NR-, -CH₂CH₂O-, -CH₂CH₂S-, -CH₂CH₂NR-, -CH₂CH₂CH₂O-, -CH₂CH₂CH₂S-, -CH₂CH₂CH₂NR-, -CH₂CH₂CH₂CH₂O-, -CH₂CH₂CH₂CH₂S-, -CH₂CH₂CH₂CH₂NR-, -CH₂CH₂OCH₂CH₂-, -(CH₂)₄NHCH₂-, -(CH₂)₃NHCH₂CH₂-, or -CH₂CH₂NHCH₂CH₂-, and preferred R' groups are hydrogen, C₁-C₄alkyl, optionally substituted tetrahydropyranyl, pyrrolidinyl, piperidinyl, piperazinyl, morpholinyl, thiomorpholinyl, pyridinyl, phenyl, or cyclohexyl, or R and R', taken together with the nitrogen atom to which they are bound, form an optionally substituted 5- or 6-membered heterocyclyl ring.

6-7. (Canceled)

8. (Previously presented) The compound of claim 1, wherein R⁶ is CH₂OH, CH₂CH₂OH, OH, OMe, OEt, NH₂, NH(Me), NH(Et), N(Me)(Me), CH₂NH₂, CH₂CH₂NH₂, NHCO₂*t*-butyl, phenyl, cyclopentyl, methyl, ethyl, isopropyl, cyclopropyl, NH(CH₂)₃NH₂, NH(CH₂)₂NH₂, NH(CH₂)₂NHEt, NHCH₂pyridyl, NHSO₂phenyl, NHC(O)CH₂C(O)O*t*-butyl, NHC(O)CH₂NH₃, and NHCH₂-imidazol-4-yl.

Applicants: Jingrong Cao et al.
Application No.: 10/696,862

9. (Previously presented) The compound of claim 3, wherein Ar¹ is:



wherein t is 0, 1, 2, 3, 4 or 5, and wherein any Ar¹ is bonded to Q² through any substitutable carbon atom, and wherein one or more hydrogen atoms on any substitutable carbon atom is substituted with one or more independent occurrences of TR⁷.

10. (Previously presented) The compound of claim 9, wherein Ar¹ is **a**, **e**, **i**, **k**, **cc**, **jj**, or **pp**.

11. (Original) The compound of claim 9, wherein T is a bond or is an optionally substituted C₁₋₆ alkylidene chain wherein one or two methylene units are optionally and

Applicants: Jingrong Cao et al.
Application No.: 10/696,862

independently replaced by -O-, -NR-, -S-, -SO₂-, -COO-, -CO-, -OSO₂-, -NRSO₂-, -CONR-, or -SO₂NR-, and R⁷ is R' or halogen.

12. (Original) The compound of claim 9, wherein each occurrence of TR⁷ is independently -C₁₋₃alkyl, -OR', -SR', -CF₃, -OCF₃, -SCF₃, -F, -Cl, I, -Br, -COOR', -COR', -O(CH₂)₄N(R)(R'), -O(CH₂)₃N(R)(R'), -O(CH₂)₂N(R)(R'), -O(CH₂)N(R)(R'), -O(CH₂)₄CON(R)(R'), -O(CH₂)₃CON(R)(R'), -O(CH₂)₂CON(R)(R'), -O(CH₂)CON(R)(R'), -C(O)N(R)(R'), -(CH₂)₄OR', -(CH₂)₃OR', -(CH₂)₂OR', -CH₂OR', optionally substituted phenyl or benzyl, -N(R)(R'), -(CH₂)₄N(R)(R'), -(CH₂)₃N(R)(R'), -(CH₂)₂N(R)(R'), -(CH₂)N(R)(R'), or SO₂N(R)(R'), NRSO₂R', CON(R)(R'), or -OSO₂R'.


13. (Canceled)

14. (Previously presented) The compound of claim 1, wherein Q³ is a direct bond, or is -(CHR⁶)_q-, -(CHR⁶)_qO-, -(CHR⁶)_qS-, -(CHR⁶)_qS(O)₂-, -(CHR⁶)_qS(O)-, -(CHR⁶)_qNR-, or -(CHR⁶)_qC(O)-, wherein q is 0, 1, 2, or 3, and R⁶ is R', -N(R)(R'), -(CH₂)₁₋₄N(R)(R'), -OR', -(CH₂)₁₋₄OR', -NR(CH₂)₁₋₄N(R)(R'), -NR(CH₂)₁₋₄SO₂R', -NR(CH₂)₁₋₄COOR', or -NR(CH₂)₁₋₄COR', or two occurrences of R⁶, taken together with the atoms to which they are bound, form an optionally substituted 3-6-membered saturated, partially unsaturated, or fully unsaturated ring.


15. (Original) The compound of claim 14, wherein R⁶ is CH₂OH, CH₂CH₂OH, OH, OMe, OEt, NH₂, NH(Me), NH(Et), N(Me)(Me), CH₂NH₂, CH₂CH₂NH₂, NHCO₂*t*-butyl, phenyl, cyclopentyl, methyl, ethyl, isopropyl, cyclopropyl, NH(CH₂)₃NH₂, NH(CH₂)₂NH₂, NH(CH₂)₂NHEt, NHCH₂pyridyl, NHSO₂phenyl, NHC(O)CH₂C(O)O*t*-butyl, NHC(O)CH₂NH₃, and NHCH₂-imidazol-4-yl.

Applicants: Jingrong Cao et al.
 Application No.: 10/696,862


16. (Previously presented) The compound of claim 1, wherein Ar² is:




a




b




c




d




e




f




g




h




i




j




k




l




m




n




o




p



q




r




s


Applicants: Jingrong Cao et al.
 Application No.: 10/696,862




t




u




v




w




x




y




z




aa




bb




cc




dd




ee




ff




gg




hh




ii



jj




kk



ll

Applicants: Jingrong Cao et al.
 Application No.: 10/696,862



wherein t is 0, 1, 2, 3, 4 or 5, and wherein any Ar^2 is bonded to Q^3 through any substitutable nitrogen or carbon atom, and wherein one or more hydrogen atoms on any substitutable nitrogen or carbon atom is substituted with one or more independent occurrences of TR^7 .

17. (Original) The compound of claim 16, wherein Ar^2 is **a, b, e, g, h, i, j, k, n, r, cc, dd, ff, jj, ll, or pp**.

18. (Original) The compound of claim 16, wherein T is a bond or is an optionally substituted C_{1-6} alkylidene chain wherein one or two methylene units are optionally and independently replaced by -O-, -NR-, -S-, - SO_2 -, -COO-, -CO-, - OSO_2 -, - NRSO_2 -, -CONR-, or - SO_2NR -, and R^7 is R' or halogen.

19. (Original) The compound of claim 16, wherein each occurrence of TR^7 is independently - $\text{C}_{1-3}\text{alkyl}$ -, -OR', -SR', - CF_3 -, - OCF_3 -, - SCF_3 -, -F-, -Cl, I-, -Br-, -COOR', -COR', - $\text{O}(\text{CH}_2)_4\text{N}(\text{R})(\text{R}')$ -, - $\text{O}(\text{CH}_2)_3\text{N}(\text{R})(\text{R}')$ -, - $\text{O}(\text{CH}_2)_2\text{N}(\text{R})(\text{R}')$ -, - $\text{O}(\text{CH}_2)\text{N}(\text{R})(\text{R}')$ -, - $\text{O}(\text{CH}_2)_4\text{CON}(\text{R})(\text{R}')$ -, - $\text{O}(\text{CH}_2)_3\text{CON}(\text{R})(\text{R}')$ -, - $\text{O}(\text{CH}_2)_2\text{CON}(\text{R})(\text{R}')$ -, - $\text{O}(\text{CH}_2)\text{CON}(\text{R})(\text{R}')$ -, -C(O)N(R)(R'), -(CH₂)₄OR', -(CH₂)₃OR', -(CH₂)₂OR', -CH₂OR', optionally substituted phenyl or benzyl, -N(R)(R'), -(CH₂)₄N(R)(R'), -(CH₂)₃N(R)(R'), -(CH₂)₂N(R)(R'), -(CH₂)N(R)(R'), or $\text{SO}_2\text{N}(\text{R})(\text{R}')$, $\text{NRSO}_2\text{R}'$, $\text{CON}(\text{R})(\text{R}')$, or - $\text{OSO}_2\text{R}'$.

Applicants: Jingrong Cao et al.
Application No.: 10/696,862

20. (Previously presented) The compound of claim 1, wherein R^5 is hydrogen, $(CH_2)_3OR'$, $(CH_2)_2OR'$, $(CH_2)OR'$, $(CH_2)_3N(R')_2$, $(CH_2)_2N(R')_2$, $(CH_2)N(R')_2$, or C_{1-4} aliphatic.

21-22. (Canceled)

23. (Original) The compound of claim 1, wherein each occurrence of R^1 is independently hydrogen, halogen, optionally substituted C_{1-4} aliphatic, OR, SR, or $N(R)_2$.

24. (Previously presented) The compound of claim 23, wherein each occurrence of R^1 is independently hydrogen, halogen, $-CH_3$, $-CH_2CH_3$, $-OH$, $-OCH_3$, $-SCH_3$, $-NH_2$, $-N(CH_3)_2$, $-N(CH_2CH_3)_2$, $-NH(CH_2)_2NHCH_3$, $-NH(cyclopropyl)$, $-NH(CH_2)cyclopropyl$, or $-NH(CH_2)_2N(CH_3)_2$.


25. (Original) The compound of claim 1, wherein each occurrence of R^Z is independently hydrogen, halogen, C_{1-4} aliphatic, OH, OR', or $N(R)(R')$.

26. (Original) The compound of claim 25, wherein each occurrence of R^Z is independently hydrogen, halogen, Me, OH, OMe, NH_2 , or $N(Me)_2$.


27. (Previously presented) The compound of claim 1, wherein R^4 groups are each independently hydrogen, C_{1-6} aliphatic, CN, $C(=O)N(R)_2$, or halogen.

Applicants: Jingrong Cao et al.
Application No.: 10/696,862

28. (Previously presented) The compound of claim 1, wherein one occurrence of R^4 is CN and compounds have the general structure **II-a**:




29. (Previously presented) The compound of claim 1, wherein R^4 is hydrogen and compounds have the general structure **III-a**:



30. (Canceled)


31. (Previously presented) The compound of claim 1, wherein R^4 is hydrogen and compounds have the general structure **VII-a**:



32. (Canceled)


Applicants: Jingrong Cao et al.
Application No.: 10/696,862

33. (Previously presented) The compound of claim 1, wherein R^4 is hydrogen and compounds have the general structure **XI-a**:




XI-a.


34. (Previously presented) The compound of claim 9, wherein Q^1 is $-\text{CO}-$, Q^2 is CHR^6 , q is 1, 2, or 3, and compounds have one of formulas **XIV**, **XV**, or **XVI**:



XIV




XV




XVI.

35. (Previously presented) The compound of claim 9, wherein Q^1 is $-\text{CO}-$, Q^2 is CHR^6 , q is 1, 2 or 3, and compounds have one of formulas **XVII**, **XVIII**, or **XIX**:


Applicants: Jingrong Cao et al.
 Application No.: 10/696,862



XVII



XVIII



XIX.

36. (Previously presented) The compound of claims 34 or 35, wherein compound variables are selected from one of more of the following groups:

- a) each occurrence of R^1 is independently hydrogen, halogen, optionally substituted C_1 - C_4 aliphatic, OR, SR, or $N(R)_2$;
- b) each occurrence of R^1 is independently hydrogen, halogen, $-CH_3$, $-CH_2CH_3$, $-OH$, $-OCH_3$, $-SCH_3$, $-NH_2$, $-N(CH_3)_2$, $-N(CH_2CH_3)_2$, $-NH(CH_2)_2NHCH_3$, $-NH(cyclopropyl)$, $-NH(CH_2)cyclopropyl$, or $-NH(CH_2)_2N(CH_3)_2$;
- c) each occurrence of R^Z is independently hydrogen, halogen, optionally substituted C_1 - C_4 aliphatic, OH, $O(R')$, or $N(R)(R')$;
- d) each occurrence of R^Z is independently hydrogen, halogen, Me, OH, OMe, NH_2 , or $N(Me)_2$;
- e) R^2 is hydrogen, or is U_nR' , where n is 1, and U is $-CH_2-$, $-CH_2CH_2-$, $-CH_2CH_2CH_2-$, $-CH_2CH_2CH_2CH_2-$, $-CH_2O-$, $-CH_2S-$, $-CH_2NR-$, $-CH_2CH_2O-$, $-CH_2CH_2S-$, $-CH_2CH_2NR-$, $-CH_2CH_2CH_2O-$, $-CH_2CH_2CH_2S-$, $-CH_2CH_2CH_2NR-$, $-CH_2CH_2CH_2CH_2O-$, $-CH_2CH_2CH_2CH_2S-$, $-CH_2CH_2CH_2CH_2NR-$, $-CH_2CH_2OCH_2CH_2-$, $-(CH_2)_4NHCH_2-$, $-(CH_2)_3NHCH_2CH_2-$, or

Applicants: Jingrong Cao et al.
Application No.: 10/696,862

-CH₂CH₂NHCH₂CH₂-, and R' groups are hydrogen, C₁-C₄alkyl, optionally substituted tetrahydropyranyl, pyrrolidinyl, piperidinyl, piperazinyl, morpholinyl, thiomorpholinyl, pyridinyl, phenyl, or cyclohexyl, or R and R', taken together with the nitrogen atom to which they are bound, form an optionally substituted 5- or 6-membered heterocyclyl ring;

f) each occurrence of R⁴ is independently hydrogen, C₁₋₆aliphatic, CN, COR, COOR, CON(R)₂, or halogen;

g) q is 1, 2, or 3;

h) R⁶ is R', -N(R)(R'), -(CH₂)₁₋₄N(R)(R'), -OR', -(CH₂)₁₋₄OR', -NR(CH₂)₁₋₄N(R)(R'), -NR(CH₂)₁₋₄SO₂R', -NR(CH₂)₁₋₄COOR', or -NR(CH₂)₁₋₄COR', or two occurrences of R⁶, taken together with the atoms to which they are bound, form an optionally substituted 3-6-membered saturated, partially unsaturated, or fully unsaturated ring;

i) R⁶ is CH₂OH, CH₂CH₂OH, OH, OMe, OEt, NH₂, NH(Me), NH(Et), N(Me)(Me), CH₂NH₂, CH₂CH₂NH₂, NHCO₂*t*-butyl, phenyl, cyclopentyl, methyl, ethyl, isopropyl, cyclopropyl, NH(CH₂)₃NH₂, NH(CH₂)₂NH₂, NH(CH₂)₂NHEt, NHCH₂pyridyl, NHSO₂phenyl, NHC(O)CH₂C(O)O*t*-butyl, NHC(O)CH₂NH₃, and NHCH₂-imidazol-4-yl;







j) Ar¹ is ring **a**, **e**, **i**, **k**, **cc**, **jj**, or **pp** wherein t is 0, 1, 2, or 3, and T is a bond or is an optionally substituted C₁₋₆ alkylidene chain wherein one or two methylene units are optionally and independently replaced by -O-, -NR-, -S-, -SO₂-, -COO-, -CO-, -OSO₂-, -NRSO₂-, -CONR-, or -SO₂NR-, and R⁷ is R' or halogen; or

k) Ar¹ is ring **a**, **e**, **i**, **k**, **cc**, **jj**, or **pp** wherein t is 0, 1, 2, or 3, and each occurrence of TR⁷ is independently -C₁₋₃alkyl, -OR', -SR', -CF₃, -OCF₃, -SCF₃, -F, -Cl, I, -Br, -COOR', -COR', -O(CH₂)₄N(R)(R'), -O(CH₂)₃N(R)(R'), -O(CH₂)₂N(R)(R'), -O(CH₂)N(R)(R'), -O(CH₂)₄CON(R)(R'), -O(CH₂)₃CON(R)(R'), -O(CH₂)₂CON(R)(R'), -O(CH₂)CON(R)(R'), -C(O)N(R)(R'), -(CH₂)₄OR', -(CH₂)₃OR', -(CH₂)₂OR', -CH₂OR',

Applicants: Jingrong Cao et al.
Application No.: 10/696,862

optionally substituted phenyl or benzyl, -N(R)(R'), -(CH₂)₄N(R)(R'), -(CH₂)₃N(R)(R'), -(CH₂)₂N(R)(R'), -(CH₂)N(R)(R'), -SO₂N(R)(R'), -NRSO₂R', -CON(R)(R'), or -OSO₂R'.

37. (Previously presented) The compound of claim 34 or 35, q is 1, and Ar¹ is optionally substituted phenyl and compounds of general formula **XIV-A** through **XIX-A** are provided:



wherein:

each occurrence of R¹ is hydrogen;

each occurrence of R^Z is hydrogen;

R² is hydrogen, or is U_nR', where n is 1, and U is -CH₂-, -CH₂CH₂-,

Applicants: Jingrong Cao et al.
Application No.: 10/696,862

-CH₂CH₂CH₂-, -CH₂CH₂CH₂CH₂-, -CH₂O-, -CH₂S-, -CH₂NR-, -CH₂CH₂O-,
-CH₂CH₂S-, -CH₂CH₂NR-, -CH₂CH₂CH₂O-, -CH₂CH₂CH₂S-, -CH₂CH₂CH₂NR-,
-CH₂CH₂CH₂CH₂O-, -CH₂CH₂CH₂CH₂S-, -CH₂CH₂CH₂CH₂NR-, -CH₂CH₂OCH₂CH₂-,
-(CH₂)₄NHCH₂-, -(CH₂)₃NHCH₂CH₂-, or -CH₂CH₂NHCH₂CH₂-, and R' groups are
hydrogen, C₁-C₄alkyl, optionally substituted tetrahydropyranyl, pyrrolidinyl, piperidinyl,
piperazinyl, morpholinyl, thiomorpholinyl, pyridinyl, phenyl, or cyclohexyl, or R and R',
taken together with the nitrogen atom to which they are bound, form an optionally
substituted 5- or 6-membered heterocyclyl ring;


each occurrence of R⁴ is independently hydrogen, C₁₋₆aliphatic, CN, CON(R)₂, or
halogen;

R⁶ is R', -N(R)(R'), -(CH₂)₁₋₄N(R)(R'), -OR', -(CH₂)₁₋₄OR',
-NR(CH₂)₁₋₄N(R)(R'), -NR(CH₂)₁₋₄SO₂R', -NR(CH₂)₁₋₄COOR', or
-NR(CH₂)₁₋₄COR'; and


t is 0, 1, 2, or 3, and each occurrence of TR⁷ is independently -C₁₋₃alkyl, -OR',
-SR', -CF₃, -OCF₃, -SCF₃, -F, -Cl, I, -Br, -COOR', -COR', -O(CH₂)₄N(R)(R'),
-O(CH₂)₃N(R)(R'), -O(CH₂)₂N(R)(R'), -O(CH₂)N(R)(R'), -O(CH₂)₄CON(R)(R'),
-O(CH₂)₃CON(R)(R'), -O(CH₂)₂CON(R)(R'), -O(CH₂)CON(R)(R'), -C(O)N(R)(R'),
-(CH₂)₄OR', -(CH₂)₃OR', -(CH₂)₂OR', -CH₂OR', optionally substituted phenyl or benzyl,
-N(R)(R'), -(CH₂)₄N(R)(R'), -(CH₂)₃N(R)(R'), -(CH₂)₂N(R)(R'),
-(CH₂)N(R)(R'), -SO₂N(R)(R'), -NRSO₂R', -CON(R)(R'), or -OSO₂R'.

38. (Previously presented) The compound of claim 16, wherein R² and Q¹-R³, taken
together with the atoms to which they are bound form a 5-membered cyclic group, and
compounds have the general formula **XX** through **XXV**:


Applicants: Jingrong Cao et al.
 Application No.: 10/696,862




XX




XXI




XXII



XXIII




XXIV




XXV.

39. (Previously presented) The compound of claim 16, R^2 and Q^1-R^3 , taken together with the atoms to which they are bound form a 5-membered cyclic group, and compounds have the general formula **XXVI** through **XXXI**:




XXVI




XXVII


Applicants: Jingrong Cao et al.
 Application No.: 10/696,862




XXVIII



XXIX




XXX




XXXI.



40. (Previously presented) The compound of claim 16, wherein R^2 and Q^1-R^3 , taken together with the atoms to which they are bound form a 6-membered cyclic group, and compounds have the general formula **XXXII** through **XXXVII**:



XXXII




XXXIII




Applicants: Jingrong Cao et al.
 Application No.: 10/696,862

XXXIV



XXXVI

XXXV



XXXVII

wherein W is O, NR⁵, or CHR⁵.

41. (Original) The compound of claims 38, 39 or 40, wherein compound variables are selected from one of more of the following groups:

- each occurrence of R¹ is independently hydrogen, halogen, optionally substituted C₁-C₄aliphatic, OR, SR, or N(R)₂;
- each occurrence of R^Z is independently hydrogen, halogen, optionally substituted C₁-C₄aliphatic, OH, OR' or N(R)(R');
- each occurrence of R⁴ is independently hydrogen, C₁₋₆aliphatic, CN, COR, COOR, CON(R)₂, or halogen;
- R⁵ is hydrogen, (CH₂)₃OR', (CH₂)₂OR', (CH₂)OR', (CH₂)₃N(R')₂, (CH₂)₂N(R')₂, (CH₂)N(R')₂, or C₁₋₄aliphatic;
- Q³ is a direct bond, or is -(CHR⁶)_q-, -(CHR⁶)_qO-, -(CHR⁶)_qS-, -(CHR⁶)_qS(O)₂-, -(CHR⁶)_qS(O)-, -(CHR⁶)_qNR-, or -(CHR⁶)_qC(O)-, wherein q is 0, 1, 2, or 3; and
- Ar² is ring **a**, **b**, **e**, **g**, **h**, **i**, **j**, **k**, **n**, **r**, **cc**, **dd**, **ff**, **jj**, **ll**, or **pp**, wherein t is 0, 1, 2, or 3, and T is a bond or is an optionally substituted C₁₋₆ alkylidene chain wherein one or two methylene units are optionally and independently replaced by -O-, -NR-, -S-, -SO₂-, -COO-, -CO-, -OSO₂-, -NRSO₂-, -CONR-, or -SO₂NR-, and R⁷ is R' or halogen.

Applicants: Jingrong Cao et al.
Application No.: 10/696,862

42. (Previously presented) The compound of claims 38, 39 or 40, wherein compound variables are selected from one or more of the following groups:

a) each occurrence of R^1 is independently hydrogen, halogen, $-CH_3$, $-CH_2CH_3$, $-OH$, $-OCH_3$, $-SCH_3$, $-NH_2$, $-N(CH_3)_2$, $-N(CH_2CH_3)_2$, $NH(CH_2)_2NHCH_3$, $NH(cyclopropyl)$, $NH(CH_2)cyclopropyl$, or $NH(CH_2)_2N(CH_3)_2$;

b) each occurrence of R^Z is independently hydrogen, halogen, Me, OH, OMe, NH_2 , or $N(Me)_2$;

c) each occurrence of R^4 is independently hydrogen, C_{1-6} aliphatic, CN, $CON(R)_2$, or halogen;


d) R^5 is hydrogen, $(CH_2)_3OR'$, $(CH_2)_2OR'$, $(CH_2)OR'$, $(CH_2)_3N(R')_2$, $(CH_2)_2N(R')_2$, $(CH_2)N(R')_2$, or C_{1-4} aliphatic;

e) Q^3 is a direct bond, or is $-(CHR^6)_q-$, $-(CHR^6)_qO-$, $-(CHR^6)_qS-$, $-(CHR^6)_qS(O)_2-$, $-(CHR^6)_qS(O)-$, $-(CHR^6)_qNR-$, or $-(CHR^6)_qC(O)-$, wherein q is 0, 1, 2, or 3; and


f) Ar^2 is ring **a**, **b**, **e**, **g**, **h**, **i**, **j**, **k**, **n**, **r**, **cc**, **dd**, **ff**, **jj**, **ll**, or **pp**, wherein t is 0, 1, 2, or 3, and each occurrence of TR^7 is independently $-C_{1-3}$ alkyl, $-OR'$, $-SR'$, $-CF_3$, $-OCF_3$, $-SCF_3$, $-F$, $-Cl$, I , $-Br$, $-COOR'$, $-COR'$, $-O(CH_2)_4N(R)(R')$, $-O(CH_2)_3N(R)(R')$, $-O(CH_2)_2N(R)(R')$, $-O(CH_2)N(R)(R')$, $-O(CH_2)_4CON(R)(R')$, $-O(CH_2)_3CON(R)(R')$, $-O(CH_2)_2CON(R)(R')$, $-O(CH_2)CON(R)(R')$, $-C(O)N(R)(R')$, $-(CH_2)_4OR'$, $-(CH_2)_3OR'$, $-(CH_2)_2OR'$, $-CH_2OR'$, optionally substituted phenyl or benzyl, $-N(R)(R')$, $-(CH_2)_4N(R)(R')$, $-(CH_2)_3N(R)(R')$, $-(CH_2)_2N(R)(R')$, $-(CH_2)N(R)(R')$, $-SO_2N(R)(R')$, $-NRSO_2R'$, $-CON(R)(R')$, or $-OSO_2R'$.

43. (Previously presented) The compound of claims 38, 39 or 40, wherein Ar^2 is optionally substituted phenyl and compounds of general formula **XX-A**, through **XXXVII** are provided:


Applicants: Jingrong Cao et al.
Application No.: 10/696,862




XX-A




XXI-A




XXII-A




XXIII-A




XXIV-A



XXV-A




XXVI-A




XXVII-A


Applicants: Jingrong Cao et al.
 Application No.: 10/696,862




XXVIII-A




XXIX-A




XXX-A




XXXI-A




XXXII-A



XXXIII-A




XXXIV-A




XXXV-A

Applicants: Jingrong Cao et al.
 Application No.: 10/696,862



XXXVI-A



XXXVII-A.

44. (Previously presented) The compound of claim 43, wherein compound variables are selected from:

each occurrence of R^1 is hydrogen;

each occurrence of R^Z is hydrogen;

each occurrence of R^4 is independently hydrogen, C_{1-6} aliphatic, CN, $CON(R)_2$, or halogen;


R^5 is hydrogen, $(CH_2)_3OR'$, $(CH_2)_2OR'$, $(CH_2)OR'$, $(CH_2)_3N(R')_2$, $(CH_2)_2N(R')_2$, $(CH_2)N(R')_2$, or C_{1-4} aliphatic;

Q^3 is a direct bond, or is $-(CHR^6)_q-$, $-(CHR^6)_qO-$, $-(CHR^6)_qS-$, $-(CHR^6)_qS(O)_2-$, $-(CHR^6)_qS(O)-$, $-(CHR^6)_qNR-$, or $-(CHR^6)_qC(O)-$, wherein q is 0, 1, 2, or 3; and


t is 0, 1, 2, or 3, and each occurrence of TR^7 is independently $-C_{1-3}$ alkyl, $-OR'$, $-SR'$, $-CF_3$, $-OCF_3$, $-SCF_3$, $-F$, $-Cl$, I , $-Br$, $-COOR'$, $-COR'$, $-O(CH_2)_4N(R)(R')$, $-O(CH_2)_3N(R)(R')$, $-O(CH_2)_2N(R)(R')$, $-O(CH_2)N(R)(R')$, $-O(CH_2)_4CON(R)(R')$, $-O(CH_2)_3CON(R)(R')$, $-O(CH_2)_2CON(R)(R')$, $-O(CH_2)CON(R)(R')$, $-C(O)N(R)(R')$, $-(CH_2)_4OR'$, $-(CH_2)_3OR'$, $-(CH_2)_2OR'$, $-CH_2OR'$, optionally substituted phenyl or benzyl, $-N(R)(R')$, $-(CH_2)_4N(R)(R')$, $-(CH_2)_3N(R)(R')$, $-(CH_2)_2N(R)(R')$, $-(CH_2)N(R)(R')$, $-SO_2N(R)(R')$, $-NRSO_2R'$, $-CON(R)(R')$, or $-OSO_2R'$.

Applicants: Jingrong Cao et al.
Application No.: 10/696,862


45. (Previously presented) The compound of claim 1, having one of the structures:




I-A-1




I-A-2




I-A-3




I-A-4




I-A-5




I-A-6




I-A-7




I-A-8




I-A-9



I-A-10




I-A-11




I-A-12


Applicants: Jingrong Cao et al.
Application No.: 10/696,862




I-A-13




I-A-14




I-A-15




I-A-16




I-A-17




I-A-18




I-A-19




I-A-20




I-A-21



I-A-22




I-A-23




I-A-24


Applicants: Jingrong Cao et al.
Application No.: 10/696,862




I-A-25




I-A-26




I-A-27




I-A-28




I-A-29




I-A-30




I-A-31




I-A-32




I-A-33



I-A-34




I-A-35




I-A-36


Applicants: Jingrong Cao et al.
Application No.: 10/696,862




I-A-37




I-A-38




I-A-39




I-A-40




I-A-41




I-A-42




I-A-43




I-B-6




I-B-19




I-B-20




I-B-21



I-B-22




I-B-23




I-B-24


Applicants: Jingrong Cao et al.
Application No.: 10/696,862




I-B-25




I-B-26




I-B-27




I-B-28




I-B-29




I-B-30




I-B-31




I-B-32




I-B-33



I-B-34




I-B-35




I-B-36


Applicants: Jingrong Cao et al.
Application No.: 10/696,862




I-B-37




I-B-38




I-B-39




I-B-40




I-B-41




I-B-42




I-B-43




I-B-44




I-B-45




I-B-46




I-B-47




I-B-48



I-B-49




I-B-50




I-B-51


Applicants: Jingrong Cao et al.
Application No.: 10/696,862




I-B-52




I-B-53




I-B-54




I-B-55




I-B-56




I-B-57




I-B-58




I-B-59




I-B-60




I-B-61




I-B-62




I-B-63



I-B-64




I-B-65




I-B-66


Applicants: Jingrong Cao et al.
Application No.: 10/696,862




I-B-67




I-B-68




I-B-69




I-B-70




I-B-71




I-B-72




I-B-73




I-B-74




I-B-75




I-B-76




I-B-77




I-B-78



I-B-79




I-B-80




I-B-81


Applicants: Jingrong Cao et al.
Application No.: 10/696,862




I-B-82




I-B-83




I-B-84




I-B-85




I-B-86




I-B-88




I-B-89




I-B-90




I-B-91




I-B-92




I-B-93



I-B-94




I-B-95




I-B-96


Applicants: Jingrong Cao et al.
Application No.: 10/696,862




I-B-97




I-B-99




I-B-100




I-B-102




I-B-103




I-B-104




I-B-105




I-B-106




I-B-107




I-B-108



I-B-109




I-B-110




I-B-111


Applicants: Jingrong Cao et al.
Application No.: 10/696,862




I-B-112




I-B-113




I-B-114




I-B-115




I-B-116




I-B-117




I-B-119




I-B-120




I-B-121




I-B-122




I-B-123



I-B-124




I-B-125




I-B-126


Applicants: Jingrong Cao et al.
Application No.: 10/696,862




I-B-127




I-B-128




I-B-129




I-B-130




I-B-146




I-B-148




I-B-149




I-B-150




I-B-151



I-B-152




I-B-153




I-B-154


Applicants: Jingrong Cao et al.
Application No.: 10/696,862




I-B-157




I-B-158




I-B-159




I-B-160




I-B-161




I-B-162




I-B-163




I-B-164




I-B-165




I-B-171




I-B-175



I-B-176




I-B-179




I-B-180


Applicants: Jingrong Cao et al.
Application No.: 10/696,862




I-B-181




I-B-182




I-B-185




I-B-186




I-B-187




I-B-190



I-B-192




I-B-193




I-B-194


Applicants: Jingrong Cao et al.
Application No.: 10/696,862




I-B-197




I-B-198




I-B-199




I-B-200




I-B-201




I-B-202




I-B-203




I-B-204




I-B-205



I-B-206




I-B-207




I-B-208


Applicants: Jingrong Cao et al.
Application No.: 10/696,862




I-B-209




I-B-210




I-B-211




I-B-212




I-B-213




I-B-214




I-B-215



I-B-216




I-B-217




I-B-218


Applicants: Jingrong Cao et al.
Application No.: 10/696,862




I-B-219




I-B-220




I-B-221




I-B-222




I-B-223




I-B-224




I-B-225



I-B-226




I-B-227




I-B-228


Applicants: Jingrong Cao et al.
Application No.: 10/696,862




I-B-229




I-B-230




I-B-231




I-B-240




I-B-241




I-B-242




I-B-243




I-B-244




I-B-245



I-B-246




I-B-247




I-B-248


Applicants: Jingrong Cao et al.
Application No.: 10/696,862




I-B-249




I-B-250




I-B-251




I-B-252




I-B-253




I-B-254




I-B-275




I-B-276




I-B-277




I-B-278



I-B-279




I-B-280




I-B-281


Applicants: Jingrong Cao et al.
Application No.: 10/696,862




I-B-282




I-B-283




I-B-284




I-B-285




I-B-286




I-B-287




I-B-288




I-B-289



I-B-290




I-B-291




I-B-292


Applicants: Jingrong Cao et al.
Application No.: 10/696,862




I-B-293




I-B-294




I-B-295




I-B-296




I-B-297




I-B-298




I-B-299




I-B-300




I-B-301




I-B-302



I-B-303




I-B-304




I-B-306


Applicants: Jingrong Cao et al.
Application No.: 10/696,862




I-B-307




I-B-308




I-B-309




I-B-312




I-B-313




I-B-314




I-B-315




I-B-321



I-B-322




I-B-323




I-B-324


Applicants: Jingrong Cao et al.
Application No.: 10/696,862




I-B-325




I-B-326




I-B-327




I-B-328




I-B-329




I-B-330




I-B-331




I-B-332




I-B-333




I-B-334




I-B-335



I-B-336




I-B-337




I-B-338


Applicants: Jingrong Cao et al.
Application No.: 10/696,862




I-B-339




I-B-340




I-B-341




I-B-342




I-B-343




I-B-344




I-B-345




I-B-346




I-C-1




I-C-2




I-C-3



I-C-4




I-C-5




I-C-6


Applicants: Jingrong Cao et al.
Application No.: 10/696,862




I-C-7




I-C-8




I-C-9




I-C-10




I-C-11




I-C-12






I-C-13



I-C-14




I-C-15




Applicants: Jingrong Cao et al.
Application No.: 10/696,862

I-C-16




I-C-19

I-C-17




I-C-20


I-C-18




I-C-21




I-C-22




I-C-23




I-C-24




I-C-25




I-C-26



I-C-27




I-C-28




I-C-30


Applicants: Jingrong Cao et al.
Application No.: 10/696,862




I-C-31




I-C-32




I-C-34




I-C-36




I-C-37



I-C-38



I-C-39



or **I-C-41**.


46. (Original) A composition comprising an effective amount of compound of claim 1, and a pharmaceutically acceptable carrier, adjuvant, or vehicle.

47-53. (Canceled)


54. (Currently amended) A method of treating or lessening the severity of a disease or disorder selected from glaucoma, Alzheimer's disease, an allergy, asthma, or diabetes in

Applicants: Jingrong Cao et al.
 Application No.: 10/696,862

a patient, said method comprising administering to said patient a compound according to claim 1 ~~or a composition comprising a compound having the formula:~~



~~or a pharmaceutically acceptable salt thereof, wherein:~~



~~R^1 is halogen, CN, NO_2 , or V_mR ;~~

~~Z^1 and Z^2 are each independently CR^Z , and Z^2 is CR^1 ;~~

~~each occurrence of R^Z is independently halogen, CN, NO_2 , or U_nR' ;~~

~~R^2 is U_nR' ;~~

~~X^1 and X^2 are each independently CR^4 or N;~~

~~each occurrence of R^4 is independently halogen, CN, NO_2 , or V_mR ;~~

~~each occurrence of U or V is independently an optionally substituted C_{1-6} alkylidene chain, wherein up to two methylene units of the chain are optionally and independently replaced by NR, S, O, CS, CO_2 , OCO, CO, COCO, CONR, NRCO, $NRCO_2$, SO_2NR , $NRSO_2$, CONRNR, NRCONR, OCONR, NRNR, $NRSO_2NR$, SO, or SO_2 ;~~

~~m and n are each independently 0 or 1;~~

~~each occurrence of R is independently hydrogen or an optionally substituted C_{1-6} aliphatic group; and each occurrence of R^1 is independently hydrogen or an optionally substituted C_{1-6} aliphatic group, a 3-8 membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully~~


Applicants: Jingrong Cao et al.
 Application No.: 10/696,862

~~unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur; or R and R², two occurrences of R, or two occurrences of R², are taken together with the atom(s) to which they are bound to form an optionally substituted 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur;~~

~~Q¹ is CO, SO₂, or SO₂NR;~~

~~R³ is Q²-Ar¹;~~

~~or R² and Q¹-R³, taken together with the nitrogen atom, form the cyclic group:~~



~~, where s is 1 or 2, each occurrence of Y is independently, as valency and stability permit, CO, CS, SO₂, O, S, NR⁵, or C(R⁵)₂, and R⁵ is U_nR';~~

~~—Q² and Q³ are each independently a bond or a C₁₋₆ alkylidene chain, wherein up to two methylene units of the chain are each optionally and independently replaced by S, O, CS, CO₂, OCO, CO, COCO, CONR', NR'CO, NR'CO₂, SO₂NR', NR'SO₂, CONR'NR', NR'CONR', OCONR', NR'NR', NR'SO₂NR', SO, or SO₂; and wherein any carbon atom in the one or more methylene units is optionally substituted with one or two occurrences of R⁶, wherein each occurrence of R⁶ is independently halogen, CN, NO₂, or U_nR', or two occurrences of R⁶, or R' and R⁶, taken together with the atoms to which they are bound, form an optionally substituted 3-6 membered cycloalkyl, heterocyclyl, aryl or heteroaryl ring;~~

~~Ar¹ is a 5-8 membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from oxygen or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from oxygen or sulfur; wherein Ar¹ is optionally substituted with 0-5 independent occurrences of TR⁷; wherein T is a~~

Applicants: Jingrong Cao et al.
Application No.: 10/696,862

~~bond or is a C₁-C₆ alkylidene chain wherein up to two methylene units of T are optionally and independently replaced by NR, S, O, CS, CO₂, OCO, CO, COCO, CONR, NRCO, NRCO₂, SO₂NR, NRSO₂, CONRNR, NRCONR, OCONR, NRNR, NRSO₂NR, SO, or SO₂;~~

~~Ar² is a 5-8 membered saturated, partially unsaturated, or fully unsaturated monocyclic ring having 0-3 heteroatoms independently selected from nitrogen, oxygen, or sulfur, or an 8-12 membered saturated, partially unsaturated, or fully unsaturated bicyclic ring system having 0-5 heteroatoms independently selected from nitrogen, oxygen, or sulfur; wherein Ar² is optionally substituted with 0-5 independent occurrences of TR⁷; wherein T is a bond or is a C₁-C₆ alkylidene chain wherein up to two methylene units of T are optionally and independently replaced by NR, S, O, CS, CO₂, OCO,~~

~~CO, COCO, CONR, NRCO, NRCO₂, SO₂NR, NRSO₂, CONRNR, NRCONR, OCONR, NRNR, NRSO₂NR, SO, or SO₂;~~

~~each occurrence of R⁷ is independently R', halogen, NO₂, or CN; and~~

~~each of the optional substituents of said aryl or heteroaryl ring is selected from halogen; R^o; OR^o; SR^o; phenyl optionally substituted with R^o; O(phenyl), optionally substituted with R^o; (CH₂)₁₋₂(phenyl), optionally substituted with R^o; CH=CH(phenyl), optionally substituted with R^o; NO₂; CN; N(R^o)₂; NR^oC(O)R^o; NR^oC(S)R^o; NR^oC(O)N(R^o)₂; NR^oC(S)N(R^o)₂; NR^oCO₂R^o; NR^oNR^oC(O)R^o; NR^oNR^oC(O)N(R^o)₂; NR^oNR^oCO₂R^o; C(O)C(O)R^o; C(O)CH₂C(O)R^o; CO₂R^o; C(O)R^o; C(S)R^o; C(O)N(R^o)₂; C(S)N(R^o)₂; OC(O)N(R^o)₂; OC(O)R^o; C(O)N(OR^o)R^o; C(NOR^o)R^o; S(O)₂R^o; S(O)₂R^o; SO₂N(R^o)₂; S(O)R^o; NR^oSO₂N(R^o)₂; NR^oSO₂R^o; N(OR^o)R^o; C(=NH)N(R^o)₂; P(O)₂R^o; PO(R^o)₂; OPO(R^o)₂; (CH₂)₀₋₂NHC(O)R^o; wherein each independent occurrence of R^o is selected from hydrogen, an optionally substituted C₁₋₆ aliphatic, an unsubstituted 5-6 membered heteroaryl or heterocyclic ring, phenyl, O(phenyl), or CH₂(phenyl), wherein optional~~

Applicants: Jingrong Cao et al.
Application No.: 10/696,862

~~substituents on the aliphatic group of R^o are selected from NH₂, NH(C₁₋₄aliphatic), N(C₁₋₄aliphatic)₂, halogen, C₁₋₄aliphatic, OH, O(C₁₋₄aliphatic), NO₂, CN, CO₂H, CO₂(C₁₋₄aliphatic), O(haloC₁₋₄aliphatic), or haloC₁₋₄aliphatic, or two independent occurrences of R^o, on the same substituent or different substituents, taken together with the atom(s) to which each R^o group is bound, form a 3-12 membered saturated, partially unsaturated, or fully unsaturated monocyclic or bicyclic ring having 0-4 heteroatoms independently selected from nitrogen, oxygen, or sulfur; and~~

~~each of the optional substituents on said alkylidene chain, aliphatic, cycloalkyl, or heterocyclyl is selected from the list of optional substituents of optional substituents for aryl and heteroaryl rings and further comprise =O, =S, =NNHR^{*}, =NN(R^{*})₂, =NNHC(O)R^{*}, =NNHCO₂(alkyl), =NNHSO₂(alkyl), or =NR^{*}, where each R^{*} is independently selected from hydrogen or a C₁₋₆ aliphatic group.~~

55. (Currently amended) The method of claim 54, wherein said method ~~compound or composition~~ is used to treat or lessen the severity of an allergy or asthma.

56. (Currently amended) The method of claim 54, wherein said method ~~compound or composition~~ is used to treat or lessen the severity of diabetes.

57. (Currently amended) The method of claim 54, wherein said method ~~compound or composition~~ is used to treat or lessen the severity of glaucoma.